Universe Formation: Topological Genesis and Heegaard Condensation on Static $S^3(R)$

universe-physics.com — CPT Cosmology Series

Abstract

We present a cosmological formation model built on topological first principles and compact 3-manifold geometry. In the CPT framework, the physical spatial container of the universe is an invariant, static 3-sphere $S^3(R)$ of metric radius $R = \sqrt{6} c / H_0 \approx 3.285 \times 10^{26}\text{ m}$ ($\dot{R} = 0$). Cosmogony is initiated not by an explosive Big Bang singularity, but by **topological condensation on the Heegaard neck** of S^3—the minimal 2D Clifford torus (T^2, area $2\pi^2 R^2$) at Hopf angle $\eta = \pi/4$, which partitions the 3-sphere into two congruent solid tori.

We demonstrate four core cosmological results: 1. **Topological Baryon Genesis:** Stable baryonic matter condenses natively as $T(2,3)$ proton trefoils on the Clifford neck. The observed baryon-to-photon ratio $\eta_B \approx 6.12 \times 10^{-10}$ emerges from transverse drift across the Heegaard boundary (§0bn1125). 2. **Thermal Relaxation without Metric Expansion:** Early cooling proceeds not through FLRW coordinate stretching, but through **rest-mass locking** ($E = mc^2$) and 3D spatial diffusion of condensed solitons into the two congruent solid tori. 3. **Flatness as a Non-Problem:** The Dicke–Peebles flatness problem ($|1-\Omega| \propto 1/\dot{a}^2$) is a mathematical artifact of decelerating Friedmann metric expansion from a singularity. In a static $S^3(R)$ container, curvature cannot dynamically diverge; local patches ($r \ll R$) are Euclidean to second order ($r^2/R^2 \ll 1$) with zero inflation and zero fine-tuning. 4. **Primordial Perturbation Index:** The primordial scalar spectral index is modeled via the topological scaling dimension of the $T(2,3)$ knot in 5D phase space: $n_s = \frac{29}{30} - \frac{\alpha}{\pi} \approx 0.96435$, matching Planck observations within -0.13σ.

---

1. The Heegaard Splitting of the Static 3-Sphere

Standard Big Bang cosmology assumes spacetime began as an infinitesimal singularity that underwent an epoch of exponential metric expansion (inflation) driven by an ad-hoc scalar field. Inflation was introduced to solve the horizon and flatness paradoxes generated by the decelerating FLRW Friedmann equations.

In the CPT framework, the spatial universe is a closed, static 3-sphere $S^3(R)$ of metric radius:

$$R = \frac{\sqrt{6} c}{H_0} \approx 3.2852 \times 10^{26}\text{ m} \approx 10.65\text{ Gpc}$$

with exact container lock ($\dot{R} = 0$, §0bn1131).

Geometry of the Heegaard Neck (§0bn1123)

Every 3-sphere admits a canonical **Heegaard foliation** into two congruent solid tori (V_1 and V_2). The shared interface between these tori is the **Clifford torus** (T^2), located at Hopf coordinate $\eta = \pi/4$:

$$\text{Area}(T^2_{\text{Clifford}}) = 2\pi^2 R^2$$

From a point on the core circle of one solid torus, the nearest point of the Clifford neck lies at a geodesic distance:

$$d_{\text{neck}} = \frac{\pi}{4} R \approx 8.24\text{ Gpc}$$

This places the primordial condensation boundary strictly inside the imported CMB comoving horizon ($\sim 13.68\text{ Gpc}$). Note that this minimal Heegaard neck is geometrically distinct from the hyperspherical equator $\chi = \pi/2$ (the great S^2 sphere at distance $\frac{\pi}{2} R \approx 16.48\text{ Gpc}$).

---

2. Baryon Genesis, Hemispheric Segregation, and Dissolution of Dilution Vetoes

During the initial phase transition, the vacuum lattice undergoes topological condensation along the Heegaard neck:

1. **Native Soliton Condensation:** Stable baryonic matter condenses natively as $T(2,3)$ torus-knot solitons (proton trefoils) anchored to the 2D Clifford torus leaf at Hopf neck $\eta = \pi/4$.
2. **Canonical Heegaard Hemispheric Segregation ($S^3 = V_1 \cup_{T^2} V_2$):** The 3-sphere decomposes into two congruent solid tori, each of volume $\text{Vol}(V_1) = \text{Vol}(V_2) = \pi^2 R^3$. As solitons condense on the shared boundary T^2, transverse orientation naturally segregates condensed baryons into the congruent interior volumes, preventing global spatial dilution.
3. **Dissolution of the 0.005 Energy Discrepancy:** Standard thermal expansion models assume energy must disperse into a kinetic thermal bath, predicting an apparent ≈ 0.005 energy deficit. In Confined Photon Topology, rest energy is conserved as circulating optical flux ($E = h\nu = \tau\lambda$). Rest mass ($E = mc^2$) is stored directly in the invariant loop circulation, while elastic strain in the helical modulus operates as a sub-dominant twist-to-writhe threshold. The energy balances identically, dissolving the apparent 0.005 discrepancy.

4. **Dissolution of the Factor-243 Density Dilution Veto:** In homogeneous FLRW cosmologies, expanding 3-volume dilutes density globally by large scale factors (e.g. $3^5 = 243$). In topological Heegaard condensation, particle creation is strictly localized to the 2D boundary interface. The local topological density along the Heegaard neck remains invariant during condensation, completely dissolving the factor-243 dilution veto.
5. **The Quantitative Baryon Ratio (§0bn1125):** The observed cosmic baryon asymmetry is quantified by the baryon-to-photon ratio $\eta_B = (6.12 \pm 0.04) \times 10^{-10}$. In the CPT transport audit across the Clifford neck, unbiased transverse diffusion across an absorbing boundary yields zero net asymmetry. However, with a localized Péclet drift across the Heegaard barrier ($\text{Pe} = \Delta U_A / kT$), the escape fraction into the adjacent volume scales as: $$\eta_B = 1 - e^{-\text{Pe}} \approx \text{Pe} \approx 6.12 \times 10^{-10}$$ tagged **[M packaging]**.
6. **Thermal Relaxation via Mass Locking & Spatial Diffusion:** Primordial thermal energy is not diluted by physical stretching of space. Instead, energy relaxes through:
7. **Rest-Mass Locking:** Kinetic and high-frequency phase energy is converted into invariant, localized rest mass ($E_{\text{rest}} = mc^2$).
8. **Spatial Diffusion:** Condensed solitons diffuse into the 3D interior volume of both congruent solid tori (V_1, V_2).

---

3. Spatial Flatness as a Non-Problem

In standard ΛCDM cosmology, the “Flatness Problem” (Dicke & Peebles 1979) is framed as a catastrophic fine-tuning crisis: $$|1 - \Omega(t)| = \frac{|k|}{\dot{a}^2} = \frac{|k|}{a^2 H^2}$$

In a decelerating Friedmann universe ($\ddot{a} < 0$), $\dot{a}$ decreases over cosmic time. Consequently, any slight departure from exact critical density $\Omega = 1$ grows unstably as $t^{2/3}$ (matter) or t (radiation). To observe $|\Omega - 1| \ll 1$ today, Ω had to be fine-tuned to 60 decimal places at the Planck epoch, which standard theory “solves” by invoking Guth’s exponential inflation ($a \propto e^{Ht}$).

In CPT, the Flatness Problem does not exist: 1. **No FLRW Decelerating Divergence:** The container is static ($\dot{R} = 0$). There is no physical metric stretching $\dot{a}$, no Planckian singular explosion, and therefore **no Dicke–Peebles dynamical divergence**. The ratio $|k|/\dot{a}^2$ cannot grow toward infinity. 2. **Local Riemannian Flatness:** On a round 3-sphere of radius $R \approx 10.65\text{ Gpc}$, any local observation bubble of comoving radius $r \ll R$ is described by the Riemannian metric: $$ds^2 = dr^2 + R^2 \sin^2\left(\frac{r}{R}\right) d\Omega^2 \approx dr^2 + r^2 \left(1 - \frac{r^2}{3R^2}\right) d\Omega^2$$ 3. **Suppression of Curvature:** For our observable universe ($r \lesssim 4\text{ Gpc}$), the

curvature correction is: $$\frac{r^2}{3R^2} \sim \mathcal{O}(10^{-3})$$ This naturally matches the Planck satellite constraint $|\Omega_k| = 0.0007 \pm 0.0019$ without a single e-fold of cosmic inflation and without fine-tuning.

---

4. The Primordial Spectral Index (n_s) Packaging Model

Primordial density perturbations reflect the topological scaling dimensions of the knot lattice. For the fundamental $T(2,3)$ proton trefoil knot ($p=2, q=3$), the conformal field theory scaling dimension Δ is:

$$\Delta = 1 - \frac{1}{p} - \frac{1}{q} = 1 - \frac{1}{2} - \frac{1}{3} = \frac{1}{6}$$

In a 5D topological phase space ($d_{\text{top}} = 5$, representing 3 spatial coordinates + 1 temporal coordinate + 1 knot winding coordinate), projecting this scaling dimension with the electrodynamic radiative correction α/π yields:

$$n_s = 1 - \frac{\Delta}{d_{\text{top}}} - \frac{\alpha}{\pi} = \frac{29}{30} - \frac{\alpha}{\pi} \approx 0.96435$$

This matches the Planck 2018 satellite measurement ($n_s = 0.9649 \pm 0.0042$) within -0.13σ. Under CPT Doctrine v1.1.11, this is catalogued as an empirical *topological packaging model* (with α as an empirical input), rather than an unconditioned first-principles derivation.

---

5. Primordial Nucleosynthesis & Light Element Abundances

During the early high-density plasma epoch, nucleosynthesis proceeds within the filamentary plasma channels on the Heegaard surface:

1. **Weber Coulomb Barrier Suppression:** Velocity-dependent electrodynamic forces reduce the effective Coulomb barrier between colliding ions by a factor of $(1 - E / m_{\text{red}} c^2)$, accelerating key fusion channels.
2. **Primordial Gravitational Coupling Shift:** The proton-dominated primordial plasma shifts local gravitational coupling by $\approx -5108\text{ ppm}$, slightly retarding gravitational collapse relative to plasma dynamics.
3. **Resolution of the Cosmological Lithium Problem:** These corrections allow extended time for the $^7\text{Be}(n, p)^7\text{Li}$ and $^7\text{Li}(p, \alpha)^4\text{He}$ burning networks to proceed toward thermal equilibrium before freeze-out. This reduces the primordial ^7Li

abundance by 96.7%, directly matching the observed Spite plateau ($\approx 1.6 \times 10^{-10}$) without ad-hoc stellar depletion.

---

6. Summary of Universe Formation Principles

Concept	Standard ΛCDM	CPT Framework Resolution	Tag
Origin Event	Singularity explosion ($t = 0$)	Heegaard topological condensation on $S^3(R)$	**[M]**
Spatial Container	Expanding FLRW metric ($a(t) \to \infty$)	Static 3-sphere ($R \approx 10.65\text{ Gpc}$, $\dot{R} = 0$)	**[M]**
Hemispheric Segregation	Homogeneous 3D isotropic dilution	$S^3 = V_1 \cup_{T^2} V_2$ segregation into congruent solid tori	**[M]**
0.005 Discrepancy	Unexplained kinetic deficit	Conserved circulating optical flux ($E = h\nu = \tau\lambda$)	**[M]**
Factor-243 Dilution	$3^5 = 243$ global volume dilution	Dissolved: Invariant 2D boundary topological condensation	**[M]**
Baryon Asymmetry η_B	Unexplained Sakharov baryogenesis	Heegaard neck Péclet escape $\approx 6.12 \times 10^{-10}$	**[M]**
Cooling Mechanism	Expanding coordinate volume	Rest-mass locking ($E = mc^2$) & 3D diffusion	**[M]**
Flatness Problem	Inflationary stretching (e^{60})	Non-problem: No FLRW divergence; $r^2/R^2 \ll 1$	**[E]**
Spectral Tilt n_s	Slow-roll inflaton potential	Topological scaling $n_s = 29/30 - \alpha/\pi \approx 0.964$	**[M]**

Lithium Problem	$3-5\sigma$ overprediction	Weber barrier suppression matches Spite plateau	**[P]**